



Evaluation of pre-PCR processing approaches for enumeration of *Salmonella enterica* in naturally contaminated animal feed

Löfström, Charlotta; Schelin, J.; Andersson, G.; Vigre, Håkan; Norling, B.; Häggblom, P.; Hoorfar, Jeffrey; Rådström, P.

Publication date:
2013

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Löfström, C., Schelin, J., Andersson, G., Vigre, H., Norling, B., Häggblom, P., Hoorfar, J., & Rådström, P. (2013). *Evaluation of pre-PCR processing approaches for enumeration of Salmonella enterica in naturally contaminated animal feed*. Poster session presented at MedVetNet Association International Scientific Conference 2013, Lyngby, Denmark.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Evaluation of pre-PCR processing approaches for enumeration of *Salmonella enterica* in naturally contaminated animal feed

C. Löfström^a, J. Schelin^b, G. Andersson^c, H. Vigre^a, B. Norling^d,
P. Häggblom^c, J. Hoorfar^a, & P. Rådström^b

^aNational Food Institute, Technical University of Denmark, Søborg; ^bApplied Microbiology, Lund University, Sweden;
^cNational Veterinary Institute (SVA), Uppsala, Sweden; ^d Quintessence Research AB, Alunda, Sweden

Aim

To evaluate 3 pre-PCR processing strategies:

- 1) flotation-qPCR (modified from [1])
- 2) MPN-PCR (modified from [2])
- 3) qualitative culture enrichment PCR [2]

for the detection and/or quantification of *Salmonella* in naturally contaminated soy bean meal.

Introduction

Animal feed might serve as a reservoir of *Salmonella* contributing to the spread into the food chain. Levels of *Salmonella* in feed samples are low, bacteria are unevenly distributed and stressed and could therefore be hard to recover using standard culture-based methods. Due to this, there is a need for accurate, sensitive, rapid and user-friendly sample preparation methods prior to molecular analyses. Moreover, to facilitate quantitative risk assessment in the feed production chain, there is a need to enumerate *Salmonella* in naturally contaminated feed.

Methods




Figure 1. Overview of the three pre-PCR processing strategies that were investigated. Bags of naturally contaminated soy bean meal (n = 15) were analyzed in parallel with the three methods.

Conclusions

- The 3 methods provide possibilities to assess the prevalence of *Salmonella* in feed, as well as the numbers of culturable, and non-culturable cells
- Differences in results could be due to non-culturable *Salmonella* and/or a heterogeneous distribution of *Salmonella* in the feed

Results

Out of the 15 bags analyzed 6, 15 and 9 were positive for *Salmonella* with flotation-qPCR, MPN-PCR and culture enrichment PCR, respectively (Table 1).

Enumeration resulted in values of 1.8×10^2 - 7.8×10^3 CFU/g (flotation-qPCR) and 0.024 to >5.2 MPN/g (MPN-PCR) (Figure 2).

Table 1. Comparison of qualitative results with the three methods for the analysis of naturally contaminated soya bean meal samples.

Method	Qualitative result (+/-) for indicated bag with soya bean meal														
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
Flotation-qPCR	+	+	-	+	-	-	+	-	-	-	-	+	-	+	-
MPN-PCR	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
Culture enrichment PCR	+	+	+	+	-	+	-	-	+	-	+	+	-	+	-




Figure 2. Comparison of enumeration results with flotation-qPCR and MPN-PCR for the analysis of naturally contaminated soy bean meal samples.

References

1. Löfström et al. (2011) IJFM 145 Suppl 1:S103-109
2. Löfström et al. (2012) Vet. Microbiol. 158(3-4):431-435.
3. Malorny et al (2004) AEM, 70(12):7046-52.

Acknowledgements

This study was financially supported by the EU project BIOTRACER (contract no. 036272) and the Swedish Research Council for Environment, Agricultural Sciences and Spatial Planning (FORMAS).